

## Orthogeriatrics—Clinical Summary Document

# Pain management in elderly fracture patients

The assessment and effective control of pain are crucial factors in the management of geriatric fracture patients to minimize physiological stress, reduce complications, and assure optimal condition for rehabilitation and recovery.

### Key points:

#### Poorly-controlled pain is common in geriatric patients

- Poorly-controlled pain in geriatric patients can lead to postoperative complications and cause functional decline.
- Standardized assessment using scales like the numerical rating scale (NRS), the verbal rating scale (VRS), or nonverbal rating scales (PAINAID) is necessary. The goals of treatment include comfort and improved function.
- In the perioperative period, opiates, early surgery, and early mobilization are the mainstays of treatment.

#### What are the consequences of poorly-controlled pain after a hip fracture?

Poorly-controlled pain has a significant impact on the clinical and functional outcomes of geriatric fracture patients and contributes to increased postoperative complications. Uncontrolled pain increases myocardial oxygen demand, promotes delirium, increases hospital length of stay, and can impair post operative recovery and rehabilitation. Undertreated or untreated pain is also associated with missed physical therapy sessions, delayed ambulation, and impaired functional recovery within the first six months following the fracture.

#### How can pain be assessed in elderly patients?

Underreporting of pain is a common problem among elderly patients. The reasons for difficulties in the assessment of pain in the elderly include:

- Communication problems (eg, hearing or language)
- Altered mental status (delirium)
- Poorly described symptoms
- Cognitive impairment

Therefore, a standardized assessment adapted to the needs of this patient group is very helpful. To avoid inadequate pain management, the use of a standardized pain assessment is important. For patients with preserved cognitive and language functions, numerical or verbal rating scales can be used. For patients with significantly impaired cognitive function, nonverbal scales (eg, PAINAID) or other clinical clues must be sought. Self-assessment can also be used in patients with mild to moderate cognitive dysfunction. Patients with severe dementia should be assessed by the use of specific indicators such as vocalization, facial expression, and body language. See Figure 1 for examples of validated pain scales.




Figure 1: Simple descriptive pain intensity scales

Assessment should be repeated throughout the day, at rest and with activity, and before and 1 hour after administration of the pain medication. It should be performed by every member of the orthogeriatric team. Pain intensity, as well as functional distress caused by the pain, must be assessed.

### How to manage pre/postoperative pain in the elderly?

In the preoperative period, intravenous opiates (morphine 2–4 mg intravenously as needed) and early surgery are the most important contributors to pain control. Local nerve blocks (femoral nerve blocks) can also be helpful in achieving rapid pain control and minimizing opiate needs. During the postoperative period, adequate pain control can often be achieved with routine scheduled administration of acetaminophen (paracetamol) and "as needed" dosing of oral opiates (starting doses: oxycodone 2.5–5 mg every 3 hours as needed; larger doses for healthy and robust patients). Doses will depend on patient comorbidities and frailty. Nonsteroidal medication should be avoided due to gastrointestinal, renal, and cardiovascular side effects.

Routine constipation prophylaxis (eg, polyethylene glycol, senna) should be given to every patient without contraindications receiving opioids. Repeat pain assessment after opioid administration to evaluate effectiveness and side effects is essential.

### Recommended reading

**Friedman SM, Mendelson DA, Kates S L, McCann RM** Geriatric co-management of proximal femur fractures: total quality management and protocol-driven care result in better outcomes for a frail patient population. *J Am Geriatr Soc.* 2008 Jul;56(7):1349–1356..

**Morrison RS, Magaziner J, Gilbert M, et al.** Relationship between pain and opioid analgesics on the development of delirium following hip fracture. *J Gerontol A Biol Sci Med Sci.* 2003 Jan;58(1):76–81.

**Parker MJ, Griffiths R, Appadu B.** Nerve blocks (subcostal, lateral cutaneous, femoral, triple, psoas) for hip fractures. *Cochrane Database Syst Rev.* 2002;(1):CD001159.

---

### Disclaimer

Production: AO Foundation, Switzerland

## **Hazards**

Great care has been taken to maintain the accuracy of the information contained in this publication. However, the publisher, and/or the distributor, and/or the editors, and/or the authors cannot be held responsible for errors or any consequences arising from the use of the information contained in this publication. Contributions published under the name of individual authors are statements and opinions solely of said authors and not of the publisher, and/or the distributor, and/or the AO Group.

The products, procedures, and therapies described in this work are hazardous and are therefore only to be applied by certified and trained medical professionals in environments specially designed for such procedures. No suggested test or procedure should be carried out unless, in the user's professional judgment, its risk is justified.

Whoever applies products, procedures, and therapies shown or described in this work will do this at their own risk. Because of rapid advances in the medical sciences, AO recommends that independent verification of diagnosis, therapies, drugs, dosages, and operation methods should be made before any action is taken.

Although all advertising material which may be inserted into the work is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement by the publisher regarding quality or value of such product or of the claims made of it by its manufacturer.

## **Legal restrictions**

This work was produced by AO Foundation, Switzerland. All rights reserved. This publication, including all parts thereof, is legally protected by copyright. Any use, exploitation or commercialization outside the narrow limits set forth by copyright legislation and the restrictions on use laid out below, without the publisher's consent, is illegal and liable to prosecution.

This applies in particular to photostat reproduction, copying, scanning or duplication of any kind, translation, preparation of microfilms, electronic data processing, and storage such as making this publication available on Intranet or Internet.

Some of the products, names, instruments, treatments, logos, designs, etc. referred to in this publication are also protected by patents and trademarks or by other intellectual property protection laws (e.g. "AO", "ASIF", "AO/ASIF", TRIANGLE/GLOBE Logo" are registered trademarks) even though specific reference to this fact is not always made in the text. Therefore, the appearance of a name, instrument, etc. without designation as proprietary is not to be construed as a representation by the publisher that it is in the public domain.

Restrictions on use: The rightful owner of an authorized copy of this work may use it for educational and research purposes only. Single images or illustrations may be copied for research or educational purposes only. The images or illustrations may not be altered in any way and need to carry the following statement of origin "Copyright by AO Foundation, Switzerland".

Copyright © 2014 by AO Foundation, Switzerland, Clavadelstrasse 8, CH-7270 Davos Platz