

Pathway for the Delirium Module of the AO Trauma Orthogeriatrics App

Katrin Singler^{1,2}, Markus Gosch², Tobias Roth³, Sacha Beck⁴

¹Institute for Biomedicine of Aging, Friedrich-Alexander-University Erlangen-Nürnberg, Germany


²Klinikum Nürnberg, Paracelsus Private Medical University, Germany

³Medical University of Innsbruck, Austria


⁴Stadtspital Waid, Zürich, Switzerland



Responsible use of phones and tablets: These devices are a potential source of infection, distraction, and patient dissatisfaction. Consult your hospital regulations before using devices in the clinical setting.



An approach to delirium in older adults with a fracture (excluding substance-withdrawal delirium)



References

Inouye SK, VanDyck CH, Alessi CA, et al. Clarifying confusion: The Confusion Assessment Method. A new method for detecting delirium. *Ann Intern Med.* 1990; 113:941–948.

Vochteloo AJ, Moerman S, van der Burg BL, et al. Delirium risk screening and haloperidol prophylaxis program in hip fracture patients is a helpful tool in identifying high-risk patients, but does not reduce the incidence of delirium. *BMC Geriatr.* 2011 Aug 11; 11:39.

Notes (for local protocols, dosages, etc)